

# Black-hole perturbation theory: The asymptotic spectrum of the prolate spin-weighted spheroidal harmonics

Shahar Hod

*The Ruppin Academic Center, Emeq Hefer 40250, Israel*  
and

*The Hadassah Institute, Jerusalem 91010, Israel*

(Dated: April 18, 2013)

Prolate spin-weighted spheroidal harmonics play a key role in black-hole perturbation theory. In particular, the highly damped quasinormal resonances of rotating Kerr black holes are closely related to the asymptotic eigenvalues of these important functions. We here present a novel and compact derivation of the asymptotic eigenvalues of the prolate spin-weighted spheroidal harmonics. Our analysis is based on a simple trick which transforms the corresponding spin-weighted spheroidal angular equation into a Schrödinger-like wave equation which is amenable to a standard WKB analysis. Our *analytical* results for the prolate asymptotic spectrum agree with previous *numerical* computations of the eigenvalues which appear in the literature.

## I. INTRODUCTION.

The characteristic dynamics of test fields in black-hole spacetimes has been studied extensively since the pioneering work of Regge and Wheeler [1], see also [2–4] and references therein. An astrophysically realistic model of wave dynamics in black-hole spacetimes should involve a non-spherical background geometry with angular momentum. In terms of the Boyer-Lindquist coordinates, the spacetime of a rotating Kerr black hole is described by the line-element [5, 6]

$$ds^2 = -\left(1 - \frac{2Mr}{\rho^2}\right)dt^2 - \frac{4Mar \sin^2 \theta}{\rho^2} dt d\phi + \frac{\rho^2}{\Delta} dr^2 + \rho^2 d\theta^2 + \left(r^2 + a^2 + \frac{2Ma^2 r \sin^2 \theta}{\rho^2}\right) \sin^2 \theta d\phi^2, \quad (1)$$

where  $M$  and  $a$  are the mass and angular momentum per unit mass of the black hole, respectively. (We use gravitational units in which  $G = c = 1$ ). Here  $\Delta \equiv r^2 - 2Mr + a^2$  and  $\rho \equiv r^2 + a^2 \cos^2 \theta$ .

In this paper we consider perturbations of the *non*-spherical Kerr spacetime. The dynamics of a test field  $\Psi$  in the rotating Kerr spacetime is governed by the well-known Teukolsky master equation [7]. One may decompose the field as

$${}_s\Psi_{lm}(t, r, \theta, \phi) = e^{im\phi} {}_sS_{lm}(\theta; a\omega) {}_s\psi_{lm}(r) e^{-i\omega t}, \quad (2)$$

where  $\omega$  is the (conserved) frequency of the mode,  $l$  is the spheroidal harmonic index, and  $m$  is the azimuthal harmonic index. The parameter  $s$  is called the spin weight of the field, and is given by  $s = \pm 2$  for gravitational perturbations,  $s = \pm 1$  for electromagnetic perturbations,  $s = \pm \frac{1}{2}$  for massless neutrino perturbations, and  $s = 0$  for scalar perturbations [7].

With the decomposition (2),  $\psi$  and  $S$  obey radial and angular equations, both of confluent Heun type [7–11], coupled by a separation constant  $A(a\omega)$ . The radial Teukolsky equation is given by [7]

$$\Delta^{-s} \frac{d}{dr} \left( \Delta^{s+1} \frac{d\psi}{dr} \right) + \left[ \frac{K^2 - 2is(r-M)K}{\Delta} - a^2\omega^2 + 2ma\omega - A + 4is\omega r \right] \psi = 0, \quad (3)$$

where  $K \equiv (r^2 + a^2)\omega - am$ .

The angular functions  $S(\theta; a\omega)$  are the spin-weighted spheroidal harmonics which are solutions of the angular equation [7–11]

$$\frac{1}{\sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{\partial S}{\partial \theta} \right) + \left[ c^2 \cos^2 \theta - 2cs \cos \theta - \frac{(m + s \cos \theta)^2}{\sin^2 \theta} + s + A \right] S = 0, \quad (4)$$

where  $c \equiv a\omega$ . The angular functions are required to be regular at the poles  $\theta = 0$  and  $\theta = \pi$ . These boundary conditions pick out a discrete set of eigenvalues  $\{{}_sA_{lm}\}$  labeled by the integers  $m$  and  $l$ . [In the  $c \rightarrow 0$  limit these angular functions become the familiar spin-weighted spherical harmonics with the corresponding angular eigenvalues  $A = l(l+1) - s(s+1) + O(a\omega)$ .]

The spin-weighted spheroidal harmonics  $S(\theta; c)$  and their corresponding eigenvalues  $\{{}_sA_{lm}\}$  have attracted much attention over the years from both physicists and mathematicians [11–17]. It is worth emphasizing that in order

to compute the characteristic resonances of black holes [2], one must *first* compute the (closely related) angular eigenvalues  $\{ {}_s A_{lm} \}$  [see Eq. (3)]. In particular, in the framework of semi-classical general relativity, it has been conjectured that the highly damped resonances may shed light on the quantum properties of black holes [18, 19]. For rotating black holes, these highly damped resonances are characterized by  $c_I \rightarrow \infty$  (where  $c_I \equiv \Im c$ ).

The asymptotic limit  $c_I \rightarrow \infty$  of the angular equation (4) was studied in [11, 12] for the  $s = 0$  case. It was found that the asymptotic scalar eigenvalues are given by

$${}_0 A_{lm} = [2(l - |m|) + 1] |c_I| + O(1) . \quad (5)$$

As pointed out in [16], the analysis of [11, 12] for the spin-0 case is somewhat incomplete. The analysis of [11, 12] requires the knowledge of the number of zeros of the scalar harmonics in the interval  $[0, \pi]$ . However, as emphasized in [16], the approximated solution found in [11, 12] for the  $c_I \rightarrow \infty$  limit is only valid in a region far from the end-points  $\theta = 0, \pi$ . Thus, the analysis of [11, 12] cannot rule out the possible existence of additional zeros of the angular eigenfunctions near the end-points. The possible omission of such zeros would lead to a wrong asymptotic behavior of the prolate (with  $c = ic_I$ ) eigenvalues, see [16] for details. In this respect the analysis of [11, 12] is not complete. One of the goals of the present paper is to present a more rigorous proof of the formula (5) for the asymptotic scalar eigenvalues.

The asymptotic spectrum of the prolate eigenvalues for the general spin case was first studied in [13]. However, as pointed out in [16, 20], the analysis of [13] for the general spin case is fundamentally flawed (see [16, 20] for details). The *numerical* results presented in Refs. [16, 20] for the general spin case provide evidence for the asymptotic behavior

$${}_s A_{lm} = \gamma |c_I| + O(1) , \quad (6)$$

where the function  $\gamma$  depends on the spin-parameter  $s$ , the azimuthal harmonic index  $m$ , and the spheroidal harmonic index  $l$  [see Eq. (21) below]. The main aim of the present paper is to provide a (simple) *analytical* proof for the prolate formula (6) in the *general* spin case.

## II. A COORDINATE TRANSFORMATION

It proves useful to introduce the coordinate  $x$  defined by [17, 21]

$$x \equiv \ln \left( \tan \left( \frac{\theta}{2} \right) \right) , \quad (7)$$

in terms of which the angular equation (4) becomes a Schrödinger-like wave equation of the form [22]

$$\frac{d^2 S}{dx^2} - US = 0 , \quad (8)$$

where

$$U(\theta) = (m + s \cos \theta)^2 - \sin^2 \theta (c^2 \cos^2 \theta - 2cs \cos \theta + s + A) . \quad (9)$$

Note that the interval  $\theta \in [0, \pi]$  maps into  $x \in [-\infty, \infty]$ . The Schrödinger-type angular equation (8) is now in a form that is amenable to a standard WKB analysis.

## III. THE SPIN-0 (SCALAR) CASE.

We shall first consider the spin-0 (scalar) case. In this case the effective potential  $U(\theta)$  is in the form of a symmetric (invariant under the transformation  $\theta \rightarrow \pi - \theta$ ) potential well: in the  $c_I \rightarrow \infty$  limit it has a local minimum at

$${}_0 \theta_{\min} = \frac{\pi}{2} \quad \text{with} \quad U({}_0 \theta_{\min}) = -A + O(1) . \quad (10)$$

Regions where  $U(\theta) < 0$  are characterized by an oscillatory behavior of the wave-function  $S$  (the ‘classically allowed regions’), while regions with  $U(\theta) > 0$  (the ‘classically forbidden regions’) are characterized by an exponential behavior (evanescent waves). The ‘classical turning points’ are characterized by  $U = 0$ . There is a pair  $\{{}_0 \theta^-, {}_0 \theta^+\}$  of such turning points (with  ${}_0 \theta^- < {}_0 \theta_{\min} < {}_0 \theta^+$ ), which in the  $c_I \rightarrow \infty$  limit are located in the near vicinity of  ${}_0 \theta_{\min}$ :

$${}_0 \theta^\pm = \frac{\pi}{2} \pm \frac{\sqrt{A}}{|c_I|} + O(c_I^{-3/2}) . \quad (11)$$

A standard textbook second-order WKB approximation for the bound-state ‘energies’ of a Schrödinger-like wave equation of the form (8) yields the well-known quantization condition [23–27]

$$\int_{x^-}^{x^+} dx \sqrt{-U(x)} = (N + \frac{1}{2})\pi \quad ; \quad N = 0, 1, 2, \dots, \quad (12)$$

where  $x^\pm$  are the turning points [with  $U(x^\pm) = 0$ ] of the potential well, and  $N$  is a non-negative integer.

Using the relation  $dx/d\theta = 1/\sin\theta$ , one can write the WKB condition (12) in the form

$$\int_{\theta^-}^{\theta^+} d\theta \frac{\sqrt{-U(\theta)}}{\sin\theta} = (N + \frac{1}{2})\pi \quad ; \quad N = 0, 1, 2, \dots. \quad (13)$$

The WKB quantization condition (13) determines the eigenvalues  $\{A\}$  of the associated spin-weighted spheroidal harmonics in the large- $c_I$  limit. The relation so obtained between the eigenvalues and the parameters  $c, m, s$  and  $N$  is rather complex and involves elliptic integrals. However, given the fact that in the  $c_I \rightarrow \infty$  limit the turning points  $\theta^\pm$  lie in the vicinity of  $\theta = \frac{\pi}{2}$  [see Eq. (11)], one can approximate the integral in (13) by [28]

$$\int_{\theta^-}^{\theta^+} d\theta \sqrt{-c_I^2(\theta - \frac{\pi}{2})^2 + A} = (N + \frac{1}{2})\pi \quad ; \quad N = 0, 1, 2, \dots. \quad (14)$$

Evaluating the integral in (14) is straightforward, and one finds

$$A(N) = (2N + 1)|c_I| + O(1) \quad ; \quad N = 0, 1, 2, \dots \quad (15)$$

for the quantized spectrum. This completes our proof for the prolate asymptotic spectrum in the scalar ( $s = 0$ ) case [29].

#### IV. THE GENERAL SPIN CASE.

We shall now consider the general spin case. In this case the effective potential  $U(\theta)$  is complex-valued. Its minimum is located at  ${}_s\theta_{\min} = \frac{\pi}{2} + \frac{is}{c_I} + O(c_I^{-2})$ , while the two turning points are located at

$${}_s\theta^\pm = \frac{\pi}{2} \pm \frac{\sqrt{A}}{|c_I|} + \frac{is}{c_I} + O(c_I^{-3/2}). \quad (16)$$

A natural generalization of the WKB analysis to the case of complex-valued potentials is provided in [30]: the WKB quantization rule is given by the standard relation

$$\int_{{}_s\theta^-}^{{}_s\theta^+} d\theta \frac{\sqrt{-U(\theta)}}{\sin\theta} = (N + \frac{1}{2})\pi \quad ; \quad N = 0, 1, 2, \dots, \quad (17)$$

which can be approximated near  $\theta = \pi/2$  by [28]

$$\int_{{}_s\theta^-}^{{}_s\theta^+} d\theta \sqrt{-c_I^2(\theta - \frac{\pi}{2})^2 - 2isc_I(\theta - \frac{\pi}{2}) + A} = (N + \frac{1}{2})\pi \quad ; \quad N = 0, 1, 2, \dots. \quad (18)$$

As emphasized in [30], the integration path between the two complex turning points  $\{{}_s\theta^-, {}_s\theta^+\}$  should be chosen such that

$$\Im\{\sqrt{-U(\theta)}\} = 0 \quad (19)$$

along the integration contour [30]. In the  $c_I \rightarrow \infty$  limit this requirement is easily fulfilled by a straight line (parallel to the real  $\theta$ -axis) which connects the two turning points (16). Substituting  $\theta = \phi - \frac{is}{c_I}$  (where  $\phi \in \mathbb{R}$  runs from  ${}_0\theta^-$  to  ${}_0\theta^+$ ) into (18) and neglecting terms of order  $O(1)$ , one finds that along the path (19) the integral (16) can be written as

$$\int_{{}_0\theta^-}^{{}_0\theta^+} d\phi \sqrt{-c_I^2(\phi - \frac{\pi}{2})^2 + A} = (N + \frac{1}{2})\pi \quad ; \quad N = 0, 1, 2, \dots. \quad (20)$$

This yields

$$A(N) = (2N + 1)|c_I| + O(1) \quad ; \quad N = 0, 1, 2, \dots \quad (21)$$

for the quantized spectrum. This completes our proof for the prolate asymptotic spectrum in the general spin case [29]. It is worth emphasizing that the *analytical* formula (21) for the prolate asymptotic spectrum agrees with the *numerical* results presented in [16, 20].

## ACKNOWLEDGMENTS

This research is supported by the Carmel Science Foundation. I thank Uri Keshet, Oded Hod, Yael Oren, Arbel M. Ongo and Ayelet B. Lata for helpful discussions.

- 
- [1] T. Regge and J. A. Wheeler, Phys. Rev. **108**, 1063 (1957).
  - [2] H. P. Nollert, Class. Quantum Grav. **16**, R159 (1999); E. Berti, V. Cardoso and A. O. Starinets, Class. Quant. Grav. **26**, 163001 (2009).
  - [3] W. H. Press, Astrophys. J. **170**, L105 (1971); V. de la Cruz, J. E. Chase and W. Israel, Phys. Rev. Lett. **24**, 423 (1970); C.V. Vishveshwara, Nature **227**, 936 (1970); M. Davis, R. Ruffini, W. H. Press and R. H. Price, Phys. Rev. Lett. **27**, 1466 (1971).
  - [4] R.H. Price, Phys. Rev. D **5**, 2419 (1972); C. Gundlach, R.H. Price, and J. Pullin, Phys. Rev. D **49**, 883 (1994); J. Bicák, Gen. Relativ. Gravitation **3**, 331 (1972); E. S. C. Ching, P. T. Leung, W. M. Suen, and K. Young, Phys. Rev. Lett. **74**, 2414 (1995); E. S. C. Ching, P. T. Leung, W. M. Suen, and K. Young, Phys. Rev. D **52**, 2118 (1995); S. Hod and T. Piran, Phys. Rev. D **58**, 024017 (1998) [arXiv:gr-qc/9712041]; S. Hod and T. Piran, Phys. Rev. D **58**, 024018 (1998) [arXiv:gr-qc/9801001]; S. Hod and T. Piran, Phys. Rev. D **58**, 044018 (1998) [arXiv:gr-qc/9801059]; S. Hod and T. Piran, Phys. Rev. D **58**, 024019 (1998) [arXiv:gr-qc/9801060]; S. Hod, Phys. Rev. D **58**, 104022 (1998) [arXiv:gr-qc/9811032]; S. Hod, Phys. Rev. D **61**, 024033 (2000) [arXiv:gr-qc/9902072]; S. Hod, Phys. Rev. D **61**, 064018 (2000) [arXiv:gr-qc/9902073]; L. Barack, Phys. Rev. D **61**, 024026 (2000); S. Hod, Phys. Rev. Lett. **84**, 10 (2000) [arXiv:gr-qc/9907096]; S. Hod, Phys. Rev. D **60**, 104053 (1999) [arXiv:gr-qc/9907044]; S. Hod, Class. Quant. Grav. **26**, 028001 (2009) [arXiv:0902.0237]; S. Hod, Class. Quant. Grav. **18**, 1311 (2001) [arXiv:gr-qc/0008001]; S. Hod, Phys. Rev. D **66**, 024001 (2002) [arXiv:gr-qc/0201017]; R. J. Gleiser, R. H. Price, and J. Pullin, Class. Quant. Grav. **25**, 072001 (2008); M. Tiglio, L. E. Kidder, and S. A. Teukolsky, Class. Quant. Grav. **25**, 105022 (2008); R. Moderski and M. Rogatko, Phys. Rev. D **77**, 124007 (2008); X. He and J. Jing, Nucl. Phys. B **755**, 313 (2006); H. Koyama and A. Tomimatsu, Phys. Rev. D **65**, 084031 (2002); B. Wang, C. Molina, and E. Abdalla, Phys. Rev. D **63**, 084001 (2001); S. Hod, Phys. Rev. Lett. **81**, 4293 (1998) [arXiv:gr-qc/9812002]; G. T. Horowitz and V. E. Hubeny, Phys. Rev. D **62**, 024027 (2000); E. W. Leaver, Proc. R. Soc. A **402**, 285 (1985); S. Hod, Phys. Rev. D **75**, 064013 (2007) [arXiv:gr-qc/0611004]; S. Hod, Class. and Quant. Grav. **24**, 4235 (2007) [arXiv:0705.2306]; A. Gruzinov, arXiv:gr-qc/0705.1725; A. Pesci, Class. Quantum Grav. **24**, 6219 (2007); S. Hod, Phys. Rev. D **78**, 084035 (2008); S. Hod, Phys. Lett. B **666**, 483 (2008) [arXiv:0810.5419]; S. Hod, Phys. Rev. D **80**, 064004 (2009) [arXiv:0909.0314]; S. Hod, Phys. Lett. A **374**, 2901 (2010) [arXiv:1006.4439]; S. Hod, Phys. Rev. D **84**, 044046 (2011) [arXiv:1109.4080]; S. Hod, Phys. Lett. B **710**, 349 (2012) [arXiv:1205.5087]; B. Mashhoon, Phys. Rev. D **31**, 290 (1985); C. J. Goebel, Astrophys. J. **172**, L95 (1972); V. Cardoso, A. S. Miranda, E. Berti, H. Witek, and V. T. Zanchin, Phys. Rev. D **79**, 064016 (2009); S. R. Dolan and A. C. Ottewill, Classical Quantum Gravity **26**, 225003 (2009); S. R. Dolan, Phys. Rev. D **82**, 104003 (2010); Y. Décanini, A. Folacci, and B. Raffaelli, Phys. Rev. D **84**, 084035 (2011).
  - [5] S. Chandrasekhar, *The Mathematical Theory of Black Holes*, (Oxford University Press, New York, 1983).
  - [6] R. P. Kerr, Phys. Rev. Lett. **11**, 237 (1963).
  - [7] S. A. Teukolsky, Phys. Rev. Lett. **29**, 1114 (1972); Astrophys. J. **185**, 635 (1973).
  - [8] A. Ronveaux, *Heun's differential equations*. (Oxford University Press, Oxford, UK, 1995).
  - [9] P. P. Fiziev, e-print arXiv:0902.1277; R. S. Borissov and P. P. Fiziev, e-print arXiv:0903.3617; P. P. Fiziev, Phys. Rev. D **80**, 124001 (2009); P. P. Fiziev, Class. Quant. Grav. **27**, 135001 (2010).
  - [10] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions* (Dover Publications, New York, 1970).
  - [11] C. Flammer, *Spheroidal Wave Functions* (Stanford University Press, Stanford, 1957).
  - [12] J. Meixner and F. W. Schäfke, *Mathieusche Funktionen und Sphäroidfunktionen mit Anwendungen auf Physikalische und Technische Probleme* (Springer-Verlag, Berlin, 1954).
  - [13] R. A. Breuer, *Gravitational Perturbation Theory and Synchrotron Radiation* (Lecture Notes in Physics, Vol. 44), (Springer, Berlin 1975).
  - [14] R. A. Breuer, M. P. Ryan Jr, and S. Waller, Proc. R. Soc. London **A358**, 71 (1977).
  - [15] M. Casals and A. C. Ottewill, Phys. Rev. D **71**, 064025 (2005).
  - [16] E. Berti, V. Cardoso, and M. Casals, Phys. Rev. D **73**, 024013 (2006); Erratum-ibid. D **73**, 109902 (2006).
  - [17] S. Hod, Phys. Lett. B **717**, 462 (2012) [arXiv:1304.0529].
  - [18] S. Hod, Phys. Rev. Lett. **81**, 4293 (1998) [arXiv:gr-qc/9812002].
  - [19] U. Keshet and S. Hod, Phys. Rev. D **76**, R061501 (2007) [arXiv:0705.1179].

- [20] E. Berti, V. Cardoso and S. Yoshida, Phys. Rev. D **69**, 124018 (2004).
- [21] H. Yang, D. A. Nichols, F. Zhang, A. Zimmerman, Z. Zhang and Y. Chen, arXiv:1207.4253.
- [22] Note that in the quantum-mechanical terminology  $-U$  stands for  $\frac{2m}{\hbar^2}(E - V)$ , where  $E, V$ , and  $m$  are the total energy, potential energy, and mass of the particle, respectively.
- [23] L. D. Landau and E. M. Lifshitz, *Quantum Mechanics*, 3rd ed. (Pergamon, New York, 1977), Chap. VII.
- [24] J. Heading, *An Introduction to Phase Integral Methods* (Wiley, New York, 1962).
- [25] C. M. Bender and S. A. Orszag, *Advanced Mathematical Methods for Scientists and Engineers* (McGraw-Hill, New York, 1978), Chap. 10.
- [26] B. F. Schutz and C. M. Will, Astrophys. J. **291**, L33 (1985); S. Iyer and C. M. Will, Phys. Rev. D **35**, 3621 (1987).
- [27] Higher order corrections to the asymptotic eigenvalues [see formula (18) below] can be obtained by using a higher-order WKB analysis [26].
- [28] Here we have used the relations  $\sin \theta = 1 - \frac{1}{2}(\theta - \frac{\pi}{2})^2 + O[(\theta - \frac{\pi}{2})^4]$  and  $\cos \theta = -(\theta - \frac{\pi}{2}) + O[(\theta - \frac{\pi}{2})^3]$ .
- [29] Note that our discrete resonance parameter  $N$  corresponds to  $l - \max(|m|, |s|)$  of [11, 12].
- [30] P. D. Miller, Physica D **152/153**, 145 (2001).